































# **Livestock Production**

## **Animal Anatomy, Physiology and**

### **Animal Health**

## **Handout 12**

### **Treating Parasites**

							
							
<b>Mange, Ear-mange</b>  <i>Sarcoptes sp.</i> <i>Psoroptes sp.</i>		•	•	•	•	•	
<b>Red mites in poultry</b>  <i>Dermanyssus avium</i>							•
<b>Ticks With 1, 2 or 3 hosts</b>  <i>Boophilus</i> <i>Amblyomma sp.</i>		•	•	•	•		
<b>Spinose Ear ticks</b>  <i>Otobius megnini</i>		•	•	•	•		
<b>Fowl ticks</b>  <i>Argas persicus</i>							•
<b>Fleas</b>  <i>Ctenocephalus sp.</i>					•	•	•
<b>Biting lice, Mallophaga</b>  <i>Damalinia bovis</i> <i>ovis</i>		•	•	•	•	•	•
<b>Poultry lice Body louse, brown louse, head louse</b>  <i>Eumerocanthus sp.</i> <i>Goniodes sp.</i> <i>Lipurus sp.</i>							•
<b>Sucking lice</b>  <i>Haematopinus sp.</i>		•	•	•	•	•	•
<b>Warble flies and grubs</b>  <i>Hypoderma sp.</i>		•	•				
<b>Human bot-flies and grubs</b>  <i>Dermatobia hominis</i>		•	•				
<b>Horse bot-flies and larvae in the horse's stomach</b>  <i>Gastrophilus intestinalis</i>				•			
<b>Sheep nasal bot-flies and larvae</b>  <i>Oestrus ovis</i>					•		
<b>Myiasis Screw worm Blowfly-Strike</b>  <i>Lucilia sp.</i> <i>Chrysomya sp.</i>			•	•	•	•	•
<b>Sheep-Keds</b>  <i>Metophagus ovinus</i>				•	•		
<b>Blood-Sucking Flies Horn fly, buffalo fly stable fly, horse fly</b>  <i>Liperosia irritans</i> <i>Stomoxys calcitrans</i> <i>Tabanus rubidus</i>		•	•	•	•	•	•